



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

The effect of Zeolite A administration on intestinal calcium absorption, blood mineral concentration and calcium excretion in urin of goat kids

Schwaller, Daniela ; Wilkens, M ; Breves, G ; Liesegang, Annette

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-73146>
Book Section

Originally published at:

Schwaller, Daniela; Wilkens, M; Breves, G; Liesegang, Annette (2012). The effect of Zeolite A administration on intestinal calcium absorption, blood mineral concentration and calcium excretion in urin of goat kids. In: Szymeczko, Roman; Iben, Christine; Burlikowska, Katarzyna; Sitkowska, Beata. 16th Congress of the Society of Veterinary and Comparative Nutrition. Bydgoszcz, Poland: s.n., 58.

ESVCN



European Society of
Veterinary & Comparative
Nutrition



Host: Department of Animal Physiology
Faculty of Animal Breeding and Biology
University of Technology and Life Sciences
in Bydgoszcz

CONGRESS PROCEEDINGS

16th Congress of the European Society of Veterinary and Comparative Nutrition

September 13-15, 2012
Bydgoszcz, Poland



Disclaimer

The editors, the referees of the Scientific Committee and the Organizing Committee have been made effort to avoid any inaccurate or misleading data, opinion or statement in this publication. The data, comments and conclusions appearing in the printed manuscripts and abstracts are the sole responsibility of the contributing authors.

Accordingly, the editors of the ESVCN Congress Proceedings and the members of the Scientific and Organizing Committees and their employees accept no responsibility or liability whatsoever for the consequences of any such inaccurate or misleading data, opinion or statement.

No part of these Proceedings may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, and recording or otherwise without prior permission from the editors.

Editors:

Roman Szymeczko¹, Christine Iben², Katarzyna Burlikowska¹ and Beata Sitkowska¹

¹ **Faculty of Animal Breeding and Biology**
University of Technology and Life Sciences
Mazowiecka 28
85-084 Bydgoszcz – Poland

² **Institute of Animal Nutrition**
University of Veterinary Medicine, Vienna
Veterinärplatz 1
1210 Vienna – Austria

Printed by Multikop Sp. Z o.o.
85-027 Bydgoszcz, ul. Jagiellońska 97

ISBN 978-83-921732-2-9

The effect of Zeolite A administration on intestinal calcium absorption, blood mineral concentration and calcium excretion in urine of goat kids

D. Schwaller¹, M. R. Wilkens², G. Breves², A. Liesegang¹

¹Institute of Animal Nutrition, Vetsuisse Faculty, University of Zurich. ²Department of Veterinary Physiology, School of Veterinary Medicine Hannover.

daniela.schwaller@uzh.ch

Introduction: Several studies on the efficacy of Zeolite A in prevention of parturient paresis in dairy cows exist (1, 2). In these studies, it was shown that Zeolite A supplementation causes a stabilized blood calcium (Ca) level around parturition. However there is no knowledge on the physiological mechanisms associated with Zeolite A. The aim of this study was to investigate whether Zeolite A has an influence on the active absorption of Ca in the intestine, the mineral status of the animals and the Ca excretion in urine.

Animals, material and methods: For this purpose, 17 goat kids (Saanen breed, 11 weeks of age) were allocated to one of three groups (one control and two treatment groups) and kept in group pens on wood shavings. All animals received hay (2nd cut) and concentrate in single boxes over three weeks. The concentrate of the treatment groups was supplemented with 1.2 g/kg body weight (Zl) respectively 1.6 g/kg body weight (Zh) Zeolite A. Blood and urine samples were periodically taken during the trial. After three weeks the animals were slaughtered and samples from intestine and kidney were collected.

Blood samples were examined for Ca, phosphorus (P) and 1,25(OH)₂D₃ and urine samples for Ca and P. The samples from intestine and kidney were analysed for genes coding for proteins which are involved in active Ca absorption in intestine and reabsorption in kidney (Transient receptor potential vanilloid (TRPV) and plasma membrane calcium ATPase (PMCA)) with PCR. For statistical evaluation multivariate analysis of variance for repeated measurements (MANOVA) was performed with group as a cofactor.

Results and discussion: There was no difference between the groups for Ca concentrations in serum. P and 1,25(OH)₂D₃ concentrations in serum were significantly decreased in both treatment groups.

TRPV and PMCA in intestine were not significantly different between all groups. However, in kidney TRPV was significantly higher in treatment groups compared to the control group. Nevertheless Ca excretion tended to be higher in treatment groups. It is described for humans, that hypophosphatemia can increase urinary excretion of Ca due to an increase in 1,25(OH)₂D₃ and therefore an increase in active Ca absorption (3). In agreement with this, in Zeolite treated cows an elevated Ca excretion in urine was observed. These authors purposed an involvement of 1,25(OH)₂D₃ (2). In contrast, in our study neither intestinal active absorption nor increased 1,25(OH)₂D₃ levels in serum could be shown. However, an unknown active mechanism in rumen (4), which seems to be independent of 1,25(OH)₂D₃ (5), might play an important role on the effect of Zeolite.

Conclusion: Zeolite A supplementation in goat kids caused no alteration in blood Ca levels, but led to hypophosphatemia and an unexpected not yet understood decrease in 1,25(OH)₂D₃ level. There is a need in further investigations on the long-term action of Zeolite especially in relation to active Ca absorption in rumen.

References:

- 1) T. Thilising-Hansen, R.J. Jørgensen, J.M.D. Enemark (2002): J Dairy Sci 85: 1855-1862
- 2) H. Grabherr, M. Spolders, M. Fürll, G. Flachowsky (2008): J Anim Physiol Anim Nutr 93: 221-236
- 3) H.W. Bleich, M.J. Moore (1979): NEJM 301: 535-541
- 4) K. Sidler-Lauff, A. Boos, M. Kraenzlin, A. Liesegang (2010): J Anim Sci 88: 3598-3610
- 5) B. Schröder, W. Goebel, K. Huber, G. Breves (2001): J Vet Med 48: 353-363